

MAPS ON POSETS, AND BLOCKERS

ANDREY O. MATVEEV

ABSTRACT. An order-theoretic generalization of Seymour relations describing the connection between the set-theoretic blocker, deletion, and contraction maps on clutters, is presented.

1. INTRODUCTION

The aim of this note is to present an order-theoretic generalization of Seymour relations [13] which describe the set-theoretic blocker, deletion, and contraction maps on clutters, see (1.2) below. Those relations are a powerful tool of discrete mathematics, see, e.g., [5, 6].

A set H is called a *blocking set* (*cover*, *system of representatives*, *transversal*) for a nonempty family $\mathcal{G} = \{G_1, \dots, G_m\}$ of nonempty subsets of a finite set if it holds $|H \cap G_k| > 0$, for each $k \in \{1, \dots, m\}$. The family of all inclusion-minimal blocking sets for $\mathcal{G}$ is called the *blocker* of $\mathcal{G}$, see, e.g., [8, Chapter 8]. We denote the blocker of $\mathcal{G}$ by $\mathcal{B}(\mathcal{G})$.

A family of subsets of a finite *ground set* S is called a *clutter* or a *Sperner family* if no set from that family contains another. The empty clutter $\emptyset$ containing no subsets of S , and the clutter $\{\hat{0}\}$ whose unique set is the empty subset $\hat{0}$ of S , are called the *trivial clutters* on S . The set-theoretic *blocker map* assigns to a nontrivial clutter its blocker, and this map alternates the trivial clutters: $\mathcal{B}(\emptyset) := \{\hat{0}\}$ and $\mathcal{B}(\{\hat{0}\}) := \emptyset$, see, e.g., [4].

Let $X \subseteq S$. The set-theoretic *deletion* ($\backslash X$), and *contraction* ($/X$) maps on clutters are defined in the following way: if $\mathcal{G}$ is a nontrivial clutter on S then the *deletion* $\mathcal{G} \backslash X$ is the family $\{G \in \mathcal{G} : |G \cap X| = 0\}$, and the *contraction* $\mathcal{G}/X$ is the family of all inclusion-minimal sets from the family $\{G - X : G \in \mathcal{G}\}$. One often says that the clutters $\mathcal{G} \backslash X$ and $\mathcal{G}/X$ are those on the ground set $S - X$. The trivial clutters do not change under the *deletion* and *contraction maps*: $\emptyset \backslash X = \emptyset/X := \emptyset$ and $\{\hat{0}\} \backslash X = \{\hat{0}\}/X := \{\hat{0}\}$.

Let $\mathcal{G}$ be a clutter on the ground set S . We have

$$\mathcal{B}(\mathcal{B}(\mathcal{G})) = \mathcal{G} , \quad (1.1)$$

see [7, 9]; given a subset $X \subseteq S$, it holds (see [13]):

$$\mathcal{B}(\mathcal{G}) \backslash X = \mathcal{B}(\mathcal{G}/X) \quad \text{and} \quad \mathcal{B}(\mathcal{G})/X = \mathcal{B}(\mathcal{G} \backslash X) . \quad (1.2)$$

Key words and phrases. Antichain, blocker, closure, clutter, contraction, deletion, lattice, poset.

2000 *Mathematics Subject Classification.* 06A06, 90C27.

2. A GENERALIZATION OF RELATIONS (1.2)

We refer the reader to [14, Chapter 3] for information and terminology in the theory of posets. See, e.g., [1, Chapter IV] on the Galois correspondence and (co)closure operators.

Theorem 2.1. *Let L be a finite poset. Let $\delta : L \rightarrow L$ be an order-preserving map, and let $\gamma : L \rightarrow L$ be an order-preserving map such that*

$$\gamma(x) \geq x, \quad (2.1)$$

for all $x \in L$. Let $\beta : L \rightarrow L$ be an order-reversing map such that

$$\beta(\beta(x)) \geq x, \quad (2.2)$$

for all $x \in L$. Either of the relations (for all $x \in L$):

$$\beta(\delta(\beta(x))) \geq \gamma(x), \quad (2.3)$$

$$\beta(\gamma(\beta(x))) \geq \delta(x) \quad (2.4)$$

implies

$$\delta(\beta(z)) \leq \beta(\gamma(z)) \leq \beta(z) \leq \gamma(\beta(z)) \leq \beta(\delta(z)), \quad (2.5)$$

for any $z \in L$. Moreover, if $\beta(\beta(x)) = x$, for all $x \in L$, then either of the equalities $\beta(\delta(\beta(x))) = \gamma(x)$ and $\beta(\gamma(\beta(x))) = \delta(x)$, for all $x \in L$, implies

$$\delta(\beta(z)) = \beta(\gamma(z)) \leq \beta(z) \leq \gamma(\beta(z)) = \beta(\delta(z)), \quad (2.6)$$

for any $z \in L$.

Proof. Relation (2.1) implies

$$\beta(\gamma(z)) \leq \beta(z)$$

because the map β is order-reversing; moreover, we have

$$\beta(z) \leq \gamma(\beta(z)).$$

We now prove implication (2.3) $\implies$ (2.5).

On the one hand, with respect to (2.2), we have $\delta(\beta(z)) \leq \beta(\delta(\beta(z)))$. On the other hand, since β is order-reversing, relation (2.3) implies $\beta(\delta(\beta(z))) \leq \beta(\gamma(z))$. We obtain

$$\delta(\beta(z)) \leq \beta(\gamma(z)). \quad (2.7)$$

Further, on the one hand, relation (2.3) implies $\gamma(\beta(z)) \leq \beta(\delta(\beta(\beta(z))))$. On the other hand, since $\beta(\beta(z)) \geq z$ by (2.2), and δ is order-preserving, and β is order-reversing, we obtain $\beta(\delta(\beta(\beta(z)))) \leq \beta(\delta(z))$. We conclude that

$$\gamma(\beta(z)) \leq \beta(\delta(z)), \quad (2.8)$$

and we are done.

We now prove implication (2.4) $\implies$ (2.5).

On the one hand, with respect to (2.4), we have $\delta(\beta(z)) \leq \beta(\gamma(\beta(\beta(z))))$. On the other hand, since β is order-reversing, and γ is order-preserving, relation (2.2) implies $\beta(\gamma(\beta(\beta(z)))) \leq \beta(\gamma(z))$. We obtain (2.7).

Further, on the one hand, relation (2.2) implies $\gamma(\beta(z)) \leq \beta(\beta(\gamma(\beta(z))))$. On the other hand, since β is order-reversing, relation (2.4) implies $\beta(\beta(\gamma(\beta(z)))) \leq \beta(\delta(z))$. We come to (2.8), and we are done.

The proof of relation (2.6) is now straightforward, with respect to the argument above. $\square$

Note that since the map β in Theorem 2.1 is order-reversing, and (2.2) holds, it is a consequence of [1, Proposition 4.36(iii)] that we have

$$\beta(\beta(\beta(x))) = \beta(x) , \quad (2.9)$$

for any $x \in L$.

To illustrate Theorem 2.1, we give a comment to (1.2). Let P be a finite bounded poset of cardinality greater than one, whose least element is denoted $\hat{0}_P$. We denote by $\mathfrak{I}(A)$ and $\mathfrak{F}(A)$ the order ideal and filter of P generated by an antichain $A \subset P$, respectively. The *atoms* of P are the elements covering $\hat{0}_P$; we denote the set of all atoms of P by P^a .

The antichains in P compose a distributive lattice, denoted $\mathfrak{A}(P)$. In the present note, the antichains are ordered in the following way: if $A', A'' \in \mathfrak{A}(P)$ then we set

$$A' \leq A'' \text{ iff } \mathfrak{F}(A') \subseteq \mathfrak{F}(A'').$$

We call the least element $\hat{0}_{\mathfrak{A}(P)}$ and greatest element $\hat{1}_{\mathfrak{A}(P)}$ of $\mathfrak{A}(P)$ the *trivial antichains* in P because, in the context of the present note, those antichains are counterparts of the trivial clutters. Here $\hat{0}_{\mathfrak{A}(P)}$ is the empty antichain in P , and $\hat{1}_{\mathfrak{A}(P)}$ is the one-element antichain $\{\hat{0}_P\}$.

- If $\{a\}$ is a nontrivial one-element antichain in P then the order-theoretic *blocker* $\mathfrak{b}(a)$ of $\{a\}$ in P is the antichain

$$\mathfrak{b}(a) := \mathfrak{I}(a) \cap P^a .$$

- If A is a nontrivial antichain in P then the order-theoretic *blocker* $\mathfrak{b}(A)$ of A in P is the following meet in $\mathfrak{A}(P)$:

$$\mathfrak{b}(A) := \bigwedge_{a \in A} \mathfrak{b}(a) . \quad (2.10)$$

- The order-theoretic *blockers* of the trivial antichains in P are:

$$\mathfrak{b}(\hat{0}_{\mathfrak{A}(P)}) := \hat{1}_{\mathfrak{A}(P)} , \quad \mathfrak{b}(\hat{1}_{\mathfrak{A}(P)}) := \hat{0}_{\mathfrak{A}(P)} .$$

See [2, 3, 10, 11, 12] on blockers in posets.

The map $\mathfrak{b} : \mathfrak{A}(P) \rightarrow \mathfrak{A}(P)$ is called the order-theoretic *blocker map* on $\mathfrak{A}(P)$. That map is order-reversing, with the property $\mathfrak{b}(\mathfrak{b}(A)) \geq A$, for all $A \in \mathfrak{A}(P)$. Equality (2.9) implies

$$\mathfrak{b}(\mathfrak{b}(\mathfrak{b}(A))) = \mathfrak{b}(A) ,$$

cf. (1.1). The posets with the property $\mathfrak{b}(\mathfrak{b}(A)) = A$, for all $A \in \mathfrak{A}(P)$, are characterized in [2].

Let $X \subseteq P^a$.

- If $\{a\}$ is a nontrivial one-element antichain in P then the order-theoretic *deletion* $\{a\} \setminus X$ and *contraction* $\{a\}/X$ of $\{a\}$ in P are the antichains

$$\begin{aligned} \{a\} \setminus X &:= \begin{cases} \{a\}, & \text{if } |\mathfrak{b}(a) \cap X| = 0, \\ \hat{0}_{\mathfrak{A}(P)}, & \text{if } |\mathfrak{b}(a) \cap X| > 0, \end{cases} \\ \{a\}/X &:= \begin{cases} \{a\}, & \text{if } |\mathfrak{b}(a) \cap X| = 0, \\ \mathfrak{b}(\mathfrak{b}(a) - X), & \text{if } |\mathfrak{b}(a) \cap X| > 0 \text{ and } \mathfrak{b}(a) \not\subseteq X, \\ \hat{1}_{\mathfrak{A}(P)}, & \text{if } \mathfrak{b}(a) \subseteq X. \end{cases} \end{aligned}$$

- If A is a nontrivial antichain in P then the order-theoretic *deletion* $A \setminus X$ and *contraction* A/X of A in P are the following joins in $\mathfrak{A}(P)$:

$$A \setminus X := \bigvee_{a \in A} (\{a\} \setminus X), \quad A/X := \bigvee_{a \in A} (\{a\}/X). \quad (2.11)$$

- The order-theoretic *deletion* and *contraction* of the trivial antichains in P are:

$$\hat{0}_{\mathfrak{A}(P)} \setminus X = \hat{0}_{\mathfrak{A}(P)}/X := \hat{0}_{\mathfrak{A}(P)}, \quad \hat{1}_{\mathfrak{A}(P)} \setminus X = \hat{1}_{\mathfrak{A}(P)}/X := \hat{1}_{\mathfrak{A}(P)}.$$

The map $(\setminus X) : \mathfrak{A}(P) \rightarrow \mathfrak{A}(P)$, $A \mapsto A \setminus X$, is called the *operator of deletion* on $\mathfrak{A}(P)$; it is a coclosure operator on $\mathfrak{A}(P)$. The map $(/X) : \mathfrak{A}(P) \rightarrow \mathfrak{A}(P)$, $A \mapsto A/X$, is called the *operator of contraction* on $\mathfrak{A}(P)$; it is a closure operator on $\mathfrak{A}(P)$ [11, Theorem 2.5].

Let the poset L from Theorem 2.1 be the lattice $\mathfrak{A}(P)$. In this context, the maps $\mathfrak{b}, (\setminus X), (/X) : \mathfrak{A}(P) \rightarrow \mathfrak{A}(P)$ are instances of the maps β, δ , and γ from Theorem 2.1, respectively. In particular, (2.3) and (2.4) read as follows:

Lemma 2.2. *For any antichain A in P , the relations*

$$\mathfrak{b}(\mathfrak{b}(A) \setminus X) \geq A/X, \quad (2.12)$$

$$\mathfrak{b}(\mathfrak{b}(A)/X) \geq A \setminus X \quad (2.13)$$

hold in $\mathfrak{A}(P)$.

Proof. There is nothing to prove if A is trivial.

Let $\{a'\}$ be a nontrivial one-element antichain in P .

1. Suppose that $|\mathfrak{b}(a') \cap X| = 0$. In this case we have

$$\mathfrak{b}(\mathfrak{b}(a') \setminus X) = \mathfrak{b}(\mathfrak{b}(a')) \geq \{a'\} = \{a'\}/X$$

and

$$\mathfrak{b}(\mathfrak{b}(a')/X) = \mathfrak{b}(\mathfrak{b}(a')) \geq \{a'\} = \{a'\} \setminus X.$$

2. Suppose that $|\mathfrak{b}(a') \cap X| > 0$ and $\mathfrak{b}(a') \not\subseteq X$. In this case we have

$$\mathfrak{b}(\mathfrak{b}(a') \setminus X) = \mathfrak{b}(\mathfrak{b}(a') - X) = \{a'\}/X$$

and

$$\mathfrak{b}(\mathfrak{b}(a')/X) = \mathfrak{b}(\hat{1}_{\mathfrak{A}(P)}) = \hat{0}_{\mathfrak{A}(P)} = \{a'\} \setminus X.$$

3. If $\mathfrak{b}(a') \subseteq X$ then we have

$$\mathfrak{b}(\mathfrak{b}(a') \setminus X) = \mathfrak{b}(\mathfrak{b}(a') - X) = \mathfrak{b}(\hat{0}_{\mathfrak{A}(P)}) = \hat{1}_{\mathfrak{A}(P)} = \{a'\}/X$$

and

$$\mathfrak{b}(\mathfrak{b}(a')/X) = \mathfrak{b}(\hat{1}_{\mathfrak{A}(P)}) = \hat{0}_{\mathfrak{A}(P)} = \{a'\} \setminus X.$$

Now, let A be an arbitrary nontrivial antichain in P . On the one hand, we by definition (2.10) have

$$\mathfrak{b}(\mathfrak{b}(A) \setminus X) = \mathfrak{b}\left(\left(\bigwedge_{a \in A} \mathfrak{b}(a)\right) \setminus X\right) \quad \text{and} \quad \mathfrak{b}(\mathfrak{b}(A)/X) = \mathfrak{b}\left(\left(\bigwedge_{a \in A} \mathfrak{b}(a)\right) / X\right)$$

in $\mathfrak{A}(P)$. On the other hand, for any element $a' \in A$, we have

$$\mathfrak{b}\left(\left(\bigwedge_{a \in A} \mathfrak{b}(a)\right) \setminus X\right) \geq \mathfrak{b}(\mathfrak{b}(a') \setminus X) \geq \{a'\}/X$$

and

$$\mathfrak{b}\left(\left(\bigwedge_{a \in A} \mathfrak{b}(a)\right) / X\right) \geq \mathfrak{b}(\mathfrak{b}(a')/X) \geq \{a'\} \setminus X$$

in $\mathfrak{A}(P)$. Definitions (2.11) now imply relations (2.12) and (2.13). $\square$

With the help of relation (2.5) and Lemma 2.2, we come to the following conclusion:

Corollary 2.3 ([11], Theorem 2.6). *For any antichain A in P , the relation*

$$\mathfrak{b}(A) \setminus X \leq \mathfrak{b}(A/X) \leq \mathfrak{b}(A) \leq \mathfrak{b}(A)/X \leq \mathfrak{b}(A \setminus X)$$

holds in $\mathfrak{A}(P)$.

REFERENCES

- [1] M. Aigner, *Combinatorial Theory*, Grundlehren der Mathematischen Wissenschaften, vol. 234, Springer (1979).
- [2] A. Björner and A. Hultman, *A note on blockers in posets*, Ann. Comb., **8** (2004), 123–131.
- [3] A. Björner, I. Peeva and J. Sidman, *Subspace arrangements defined by products of linear forms*, J. London Math. Soc. (2), *to appear*.
- [4] R. Cordovil, K. Fukuda and M.L. Moreira, *Clutters and matroids*, Discrete Math., **89** (1991) 161–171.
- [5] G. Cornuéjols, *Combinatorial Optimization. Packing and Covering*, CBMS-NSF Regional Conference Series in Applied Mathematics, vol. 74, Philadelphia PA: SIAM (2001).
- [6] Y. Crama and P.L. Hammer, with contributions by C. Benzaken, J.C. Bioch, E. Boros, N. Brauner, V. Gurvich, L. Hellerstein, T. Ibaraki, A. Kogan, K. Makino, R. Pöschel, I. Rosenberg, B. Simeone and B. Vettier, *Boolean Functions*, in preparation.

- [7] J. Edmonds and D.R. Fulkerson, *Bottleneck extrema*, J. Combinatorial Theory, **8** (1970) 299–306.
- [8] M. Grötschel, L. Lovász and A. Schrijver, *Geometric Algorithms and Combinatorial Optimization* Algorithms and Combinatorics, vol. 2, Springer (1993).
- [9] A. Lehman, *A solution of the Shannon switching game*, J. Soc. Indust. Appl. Math., **12** (1964) 687–725.
- [10] A.O. Matveev, *On blockers in bounded posets*, Int. J. Math. Math. Sci., **26** (2001), no. 10, 581–588.
- [11] A.O. Matveev, *A note on operators of deletion and contraction for antichains*, Int. J. Math. Math. Sci., **31** (2002), no. 12, 725–730.
- [12] A.O. Matveev, *Extended blocker, deletion, and contraction maps on antichains*, Int. J. Math. Math. Sci., **2003** (2003), no. 10, 607–616.
- [13] P.D. Seymour, *The forbidden minors of binary clutters*, J. London Math. Soc. (2) **12** (1975/1976) 356–360.
- [14] R.P. Stanley, *Enumerative Combinatorics, vol. 1*, Cambridge Studies in Advanced Mathematics, vol 49, Cambridge University Press (1997).

DATA-CENTER CO., RU-620034, EKATERINBURG, P.O. BOX 5, RUSSIAN FEDERATION
E-mail address: aomatveev@{dc.ru, hotmail.com}